

III INTERNATIONAL SYMPOSIUM ON GENETICS AND PLANT BREEDING

OVERCOMING ABIOTIC AND BIOTIC STRESS CONSTRAINTS IN PLANT SCIENCE

CHARACTERIZATION OF LONG NON-CODING RNAS (LNCRNAS) IN MAIZE SUBJECTED TO COLONIZATION BY THE ENDOPHYTIC DIAZOTROPHIC BACTERIA HERBASPIRILLUM SEROPEDICAE AND ADDITION OF HUMIC ACID

III Simpósio Internacional de Atualização em Genética e Melhoramento de Plantas, 0ª edição, de 24/05/2021 a 26/05/2021 ISBN dos Anais: 000

> SILVA; Leandro de Oliveira ¹, VIEIRA; Lucas Maciel ², WALTER; Maria Emilia M. T. ³, VENÂNCIO; Thiago M.⁴, OLIVARES; Fábio Lopes ⁵, GRATIVOL; Clicia ⁶

RESUMO

Maize (Zea mays) is one of the most important cereal crops, used in human food, production of animal feed and bioenergy. To optimize your cultivation, it is possible to use diazotrophic and growth-promoting bacteria. However, the complex beneficial plant-bacterial relationship has not yet been elucidated, which includes the role of long non-coding RNAs (IncRNAs). The present work aims to characterize IncRNAs expressed in maize submitted to colonization by the diazotrophic bacteria Herbaspirillum seropedicae with the addition of humic acid. Reads of RNAseq libraries were aligned to the Z. mays genome by the use of HISAT2, and StringTie was used to assemble the transcripts. The differential expression of IncRNAs was calculated using DESeq2. A model based on machine learning was created to increase the accuracy of predicting new IncRNAs in maize. The search of IncRNAs in maize returned 124,739 different known IncRNAs. Through BLASTN with the transcriptome, 32,645 known IncRNAs were identified. Of these, 3,263 IncRNAs were differentially expressed (1,502 up-regulated and 1,761 down-regulated). The assembled transcripts were submitted to different tools, as a way to identify the new IncRNAs: size filter; calculation of the protein coding potential (CPC); calculation of the size of the ORF; alignment against a Rfam database and, finally, analysis with the model. A total of 6,736 new highly confidence IncRNAs were identified, 1,050 of which are differentially expressed (480 up-regulated and 570 down-regulated). The next steps include: prediction of the target genes of IncRNAs and characterization using Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG); verification of alternative splicing mechanisms; mapping of IncRNAs in public miRNA libraries and validating their targets through degradoma libraries; construction of a network of maize IncRNA-miRNA interaction. This work can contribute to a greater understanding of the roles played by maize lcnRNAs.

¹ Universidade Estadual do Norte Fluminense Darcy Ribeiro, oliveiradasilvaleo@gmail.com

- Universidade de Brasília, lucas.maciel.vieira@gmail.com
- ³ Universidade de Brasília, memtwalter@gmail.com
 ⁴ Universidade Estadual do Norte Fluminense Darcy Ribeiro, thiago.venancio@gmail.com
- Universidade Estadual do Norte Fluminense Darcy Ribeiro, fabioliv@uenf.b
- ⁶ Universidade Estadual do Norte Fluminense Darcy Ribeiro, cgrativol@uenf.br

PALAVRAS-CHAVE: Zea mays, non-coding sequences, diazotrophic bacteria

¹ Universidade Estadual do Norte Fluminense Darcy Ribeiro, oliveiradasilvaleo@gmail.com
 ² Universidade de Brasília, lucas.maciel.vieira@gmail.com
 ³ Universidade de Brasília, memtwalter@gmail.com
 ⁴ Universidade Estadual do Norte Fluminense Darcy Ribeiro, thiago.venancio@gmail.com
 ⁵ Universidade Estadual do Norte Fluminense Darcy Ribeiro, fabioliv@uenf.br
 ⁶ Universidade Estadual do Norte Fluminense Darcy Ribeiro, cgrativol@uenf.br