

METHANE EMISSIONS AT WATER-SATURATED AND DRAINED SOILS OF THE POLYGONAL TUNDRA OF SAMOYLOV ISLAND, NORTHEASTERN SIBERIA

VI Simpósio APECS-Brasil, 1ª edição, de 02/02/2021 a 04/02/2021
ISBN dos Anais: 978-65-86861-75-4

GALERA; Leonardo de Aro¹, KNOBLAUCH; Christian², ECKHARDT; Tim³, BEER; Christian⁴, PFEIFFER; Eva-Maria⁵

RESUMO

The increase in permafrost temperatures promotes thawing and release of frozen organic carbon. This leads to additional emissions of CO₂ and CH₄. It is vital to assess the amount of C that is mineralized as CH₄ due to its higher global warming potential (GWP) compared to CO₂. The most important environmental source of CH₄ is soil organic matter decomposition in anoxic conditions. Recent publications acknowledge the open debate over the strength of the permafrost carbon-climate feedback of water-saturated and drained soils. Vegetation is also an important control for CH₄ emissions due to its influence in CH₄ oxidation. Our objective was to assess CH₄ emissions in the Siberian tundra during the growing season. We performed *in situ* chamber measurements in a polygon containing two sites with distinct hydrological features in Samoylov Island in the Lena River Delta, Northeastern Siberia. The polygon center presented water-saturated conditions, while the polygon rim was well-drained. A trenching experiment allowed the quantification of the plant-mediated CH₄ transport, through the comparison of plots that had the vegetation removed and intact plots. The median CH₄ flux at the polygon center was of 26 mg.m⁻².d⁻¹, and at the polygon rim was of 1.8 mg.m⁻².d⁻¹. The CH₄ fluxes from the center varied greatly throughout the measurement period, presenting evident seasonality. The opposite was found for the rim fluxes. The polygon center median CH₄ flux decreased by 80% when the vegetation was removed, showing the importance of plant-mediated CH₄ transport in this environment. At the polygon rim there was virtually no difference between clipped and vegetated plots. The CH₄ emissions measured in this study are lower than previous studies in the same island and other permafrost-affected environments. It should be noticed that the measurements occurred only during the growing season. The off-season and winter CH₄ emissions might be significant, as has already been showed in year-around measurement studies. Our results emphasize the great pedon-scale variability of CH₄ emissions in the Siberian tundra, specially associated with hydrology, topography, and vegetation.

PALAVRAS-CHAVE: CH₄, arctic, oxic, anoxic, clipping, greenhouse gases, carbon

¹ Universität Hamburg, leonardo.galera@uni-hamburg.de

² Universität Hamburg, christian.knoblauch@uni-hamburg.de

³ Universität Hamburg, tim.eckhardt@uni-hamburg.de

⁴ Universität Hamburg, christian.beer@uni-hamburg.de

⁵ Universität Hamburg, eva-maria.pfeiffer@uni-hamburg.de